

WEST Search History

DATE: Thursday, September 15, 2005

| <u>Hide?</u> | <u>Set Name</u> | <u>Query</u> | <u>Hit Count</u> |
|--------------------------|----------------------------------|-----------------------------------|------------------|
| | <i>DB=USPT; PLUR=YES; OP=ADJ</i> | | |
| <input type="checkbox"/> | L23 | L22 and l20 | 63 |
| <input type="checkbox"/> | L22 | l19.ti,ab,clm. | 576 |
| <input type="checkbox"/> | L21 | L19 with l1 not l17 | 191 |
| <input type="checkbox"/> | L20 | L19 same l1 not l17 | 619 |
| <input type="checkbox"/> | L19 | botulin\$ or tetan\$ | 6116 |
| <input type="checkbox"/> | L18 | l17 not l11 | 130 |
| <input type="checkbox"/> | L17 | L16 with l1 | 133 |
| <input type="checkbox"/> | L16 | clostrid\$ or neurotoxin | 7518 |
| <input type="checkbox"/> | L15 | L14 same l13 same l9 not l10 | 5 |
| <input type="checkbox"/> | L14 | l4 same (l5 or l6) | 17929 |
| <input type="checkbox"/> | L13 | l1 same L12 | 5561 |
| <input type="checkbox"/> | L12 | "single chain" | 12502 |
| <input type="checkbox"/> | L11 | l3 and l10 | 35 |
| <input type="checkbox"/> | L10 | l4 same l5 same l6 same L9 | 45 |
| <input type="checkbox"/> | L9 | cleav\$ or protease or proteinase | 108341 |
| <input type="checkbox"/> | L6 | endocyt\$6 | 4351 |
| <input type="checkbox"/> | L5 | transport\$ | 460448 |
| <input type="checkbox"/> | L4 | bind\$4 | 366616 |
| <input type="checkbox"/> | L3 | l1 with L2 | 31832 |
| <input type="checkbox"/> | L2 | gene or plasmid or protein | 191203 |
| <input type="checkbox"/> | L1 | fus\$4 or chimers\$3 | 261774 |

END OF SEARCH HISTORY

File 155:MEDLINE(R) 1951-2005/Sep 19 (c) format only
2005 Dialog

| Set | Items | Description |
|-----|-------|--|
| S1 | 6955 | DC='B3.300.390.400.200.' (CLOSTRIDIUM) |
| S2 | 11321 | DC='D12.776.828.' (RECOMBINANT PROTEINS) |
| S3 | 193 | S1 AND S2 |
| S4 | 7486 | DC='D24.185.926.640.' (NEUROTOXINS) |
| S5 | 4423 | DC='D24.185.926.123.179.' (BOTULINUM TOXINS) |
| S6 | 90 | S2 AND S5 |
| S7 | 37269 | 'MUTAGENESIS, SITE-DIRECTED' |
| S8 | 18657 | DC='G5.600.' (MUTAGENESIS) |
| S9 | 28 | S5 AND S7 |
| S10 | 5 | S5 AND S8 NOT S9 |
| S11 | 13 | S5 AND PRECURSOR |
| S12 | 136 | S4 AND PRECURSOR NOT S11 |
| S13 | 45 | S7 AND S4 NOT S9 |
| S14 | 136 | S12 NOT S13 |

| Ref | Items | Index-term |
|-----|-------|-------------------------|
| E1 | 4 | AU=DOLLWET H H |
| E2 | 1 | AU=DOLLWET-MACK SUSANNE |
| E3 | 0 | *AU=DOLLY |
| E4 | 1 | AU=DOLLY C H |
| E5 | 7 | AU=DOLLY F R |
| E6 | 1 | AU=DOLLY J |
| E7 | 149 | AU=DOLLY J O |
| E8 | 9 | AU=DOLLY J OLIVER |
| E9 | 1 | AU=DOLLY JOHN P |
| E10 | 1 | AU=DOLLY M C |
| E11 | 3 | AU=DOLLY O |
| E12 | 3 | AU=DOLLY O J |
| E13 | 1 | AU=DOLLY OLIVER |
| E14 | 2 | AU=DOLLY R C |
| E15 | 1 | AU=DOLMA L |
| E16 | 1 | AU=DOLMA SONAM |

| | | |
|-----|--------|-------------------|
| S15 | 162 | E6-E8, E12 |
| S16 | 48 | S15 AND S5 |
| S17 | 0 | S7 AND S16 |
| S18 | 159426 | SUBSTITUT? |
| S19 | 2 | S18 AND S16 |
| S20 | 0 | S16 AND PRECURSOR |

6/6/1 18317639 PMID: 15721769
Preparation of specifically activatable endopeptidase toxins type A, B, and C and their applications. Mar 2005

6/6/2 18119249 PMID: 15769746
Lipid raft association of SNARE proteins regulates exocytosis in PC12 cells. May 20 2005

6/6/3 17692264 PMID: 15781237
N-terminal helix reorienters in recombinant C-fragment of Clostridium botulinum type B. Apr 29 2005

6/6/4 17726857 PMID: 15677772

Simvastatin inhibits growth factor expression and modulates profibrogenic markers in lung fibroblasts. Apr 2005

6/6/5 17561966 PMID: 15649138
Endocytosis and retrograde axonal traffic in motor neurons. 2005

6/6/6 17279925 PMID: 15359569
Use of biophysical characterization in preformulation development of a heavy-chain fragment of botulinum serotype B: evaluation of suitable purification process conditions. Aug 2004

6/6/7 16158501 PMID: 15465919
Using fluorescent sensors to detect botulinum neurotoxin activity in vitro and in living cells. Oct 12 2004

6/6/8 15414141 PMID: 15123599
Synaptotagmins I and II act as nerve cell receptors for botulinum neurotoxin G. Jul 16 2004

6/6/9 15382821 PMID: 15198662
Differential effects of Rho GTPases on axonal and dendritic development in hippocampal neurones. Jul 2004

6/6/10 15324488 PMID: 14978027
RaA-exocyst interaction mediates GTP-dependent exocytosis. May 7 2004

6/6/11 15226380 PMID: 14982988
Plasma membrane localization signals in the light chain of botulinum neurotoxin. Mar 2 2004

6/6/12 15191650 PMID: 14766296
Cloning, high-level expression, single-step purification, and binding activity of His6-tagged recombinant type B botulinum neurotoxin heavy chain transmembrane and binding domain. Mar 2004

6/6/13 15160761 PMID: 14731268
The HCC-domain of botulinum neurotoxins A and B exhibits a singular ganglioside binding site displaying serotype specific carbohydrate interaction. Feb 2004

6/6/14 15119607 PMID: 14680933
Scale-up of the fermentation and purification of the recombinant heavy chain fragment C of botulinum neurotoxin serotype F, expressed in Pichia pastoris. Nov 2003

6/6/15 15060107 PMID: 14573702
Botulinum toxin type B micromechanosensor. Nov 11 2003

6/6/16 14897423 PMID: 12750364
Entrapment of Rho ADP-ribosylated by Clostridium botulinum C3 exoenzyme in the Rho-guanine nucleotide dissociation inhibitor-1 complex. Aug 1 2003

6/6/17 14652843 PMID: 12574487
Functional interaction of auxiliary subunits and synaptic proteins with Ca(v)1.3 may impart hair cell Ca2+ current properties. Feb 2003

6/6/18 14627022 PMID: 12527421
Sequence of the gene for Clostridium botulinum type B neurotoxin associated with infant botulism, expression of the C-terminal half of heavy chain and its binding activity. Jan 3 2003

6/6/19 14493014 PMID: 12438137
Annexin 7: a non-SNARE proteolytic substrate for botulinum toxin type C in secreting chromaffin cells. Oct 2002

6/6/20 14458762 PMID: 12388760
Tenascin-C modulates matrix contraction via focal adhesion kinase- and Rho-mediated signaling pathways. Oct 2002

6/6/21 14410629 PMID: 12270602
Recovery of intracellular recombinant proteins from the yeast Pichia pastoris by cell permeabilization. Oct 23 2002

6/6/22 14404396 PMID: 12240638
Botulinum beatoen. Sep 2002

6/6/23 14397869 PMID: 12105193

Inhibition of release of neurotransmitters from rat dorsal root ganglia by a novel conjugate of a Clostridium botulinum toxin A endopeptidase fragment and Erythrina cristagalli lectin. Sep 20 2002

6/6/24 14376229 PMID: 12091381
Characterization of new cell permeable C3-like proteins that inactivate Rho and stimulate neurite outgrowth on inhibitory substrates. Sep 6 2002

6/6/25 14361569 PMID: 12177434
Potent neutralization of botulinum neurotoxin by recombinant oligoclonal antibody. Aug 20 2002

6/6/26 14341742 PMID: 12165316
Clostridium botulinum C2 toxin: binding studies with fluorescence-activated cytometry. Aug 2002

6/6/27 14257813 PMID: 12064550
Initial purification of recombinant botulinum neurotoxin fragments for pharmaceutical production using hydrophobic charge induction chromatography. Apr 5 2002

6/6/28 14177096 PMID: 11963021
New drugs 2002. Part II. Apr 2002

6/6/29 14176067 PMID: 11847234
Interaction of the Rho-ADP-ribosylating C3 exoenzyme with RaA. Apr 26 2002

6/6/30 13946422 PMID: 11700044
Site-directed mutagenesis identifies active-site residues of the light chain of botulinum neurotoxin type A. Nov 16 2001

6/6/31 13800583 PMID: 11373287
A discontinuous SNAP-25 C-terminal coil supports exocytosis. Jul 27 2001

6/6/32 13690267 PMID: 11334406
Increased intracellular calcium is required for spreading of rat islet beta-cells on extracellular matrix. May 2001

6/6/33 13687573 PMID: 11329292
Thermal stabilization of the catalytic domain of botulinum neurotoxin E by phosphorylation of a single tyrosine residue. Feb 20 2001

6/6/34 13686329 PMID: 11330351
Peptide phage display library as source for inhibitors of clostridial neurotoxins. Jan 2001

6/6/35 13595682 PMID: 11176890
From the Food and Drug Administration. Feb 14 2001

6/6/36 13451445 PMID: 10409113
Protein kinase C inhibits Kv1.1 potassium channel function. Jul 1999

6/6/37 13449522 PMID: 10413679
Functional characterisation of tetanus and botulinum neurotoxins binding domains. Aug 1999

6/6/38 13207684 PMID: 11307955
Recombinant derivatives of clostridial neurotoxins as delivery vehicles for proteins and small organic molecules. Nov 2000

6/6/39 13153741 PMID: 11186457
[Effective expression of fragments of a botulinum neurotoxin type A gene coding for the L-chain and H-chain in E. coli, with formation of products causing protective immunity to administration of the toxin] Efektivnaya ekspressiya v kleitkakh E. coli fragmentov gena botulinicheskogo neirotoksina tipa A, kodiruiushchikh L-sep' i H-sep', s obrazovaniem produktov, vyzyvaiushchikh protektivnyi immunitet k vvedeniiu toksina. 2000

6/6/40 12987028 PMID: 10945449
Recombinant antibodies: a new reagent for biological agent detection. Jan 2000

6/6/41 12899885 PMID: 10844004
Syntactin modulation of slow inactivation of N-type calcium channels. Jun 15 2000

6/6/42 12883746 PMID: 10748216
Recognition of RhoA by Clostridium botulinum C3 exoenzyme. Jun 2 2000

- 6/6/43 12874837 PMID: 10816419
Tyrosine phosphorylation of the vascular endothelial-growth-factor receptor-2 (VEGFR-2) is modulated by Rho proteins. Jun 1 2000
- 6/6/44 12802814 PMID: 10733887
Fermentation, purification, and efficacy of a recombinant vaccine candidate against botulinum neurotoxin type F from *Pichia pastoris*. Apr 2000
- 6/6/45 12724286 PMID: 10548557
nSec1 binds a closed conformation of syntaxin1A. Jan 24 2000
- 6/6/46 12721691 PMID: 10544682
Binding of *Clostridium botulinum* C2 toxin to asparagine-linked complex and hybrid carbohydrates. Jan 28 2000
- 6/6/47 12718036 PMID: 10540419
Regulation of LPA-promoted myofibroblast contraction: role of Rho, myosin light chain kinase, and myosin light chain phosphatase. Feb 1 2000
- 6/6/48 12482093 PMID: 9792657
Characterization of the catalytic site of the ADP-ribosyltransferase *Clostridium botulinum* C2 toxin by site-directed mutagenesis. Nov 6 1998
- 6/6/49 12469820 PMID: 9782457
On the action of botulinum neurotoxins A and E at cholinergic terminals. Apr 1998
- 6/6/50 12442605 PMID: 9746583
Characterization of *Clostridium botulinum* type B neurotoxin associated with infant botulism in Japan. Oct 1998
- 6/6/51 12403522 PMID: 9712688
Ganglioside GT1b as a complementary receptor component for *Clostridium botulinum* neurotoxins. Aug 1998
- 6/6/52 12383413 PMID: 9693060
Production and purification of the heavy-chain fragment C of botulinum neurotoxin, serotype B, expressed in the methylobacterial yeast *Pichia pastoris*. Aug 1998
- 6/6/53 12305846 PMID: 9618155
Regulation of TNF- α -induced reorganization of the actin cytoskeleton and cell-cell junctions by Rho, Rac, and Cdc42 in human endothelial cells. Jul 1998
- 6/6/54 12195318 PMID: 9494111
Guanine nucleotide-dependent translocation of RhoA from cytosol to high affinity membrane binding sites in human erythrocytes. Mar 15 1998
- 6/6/55 12182086 PMID: 9478920
The alpha-subunit of the heterotrimeric G protein G13 activates a phosphatase D isozyme by a pathway requiring Rho family GTPases. Feb 27 1998
- 6/6/56 12130139 PMID: 9430593
Postsynaptic membrane fusion and long-term potentiation. Jan 16 1998
- 6/6/57 12114293 PMID: 9414082
Botulinum neurotoxin types A and E require the SNARE motif in SNAP-25 for proteolysis. Nov 24 1997
- 6/6/58 12094228 PMID: 9393976
Inactivation of the small GTPase Rho disrupts cellular attachment and induces adhesion-dependent and adhesion-independent apoptosis. Oct 1997
- 6/6/59 12056776 PMID: 9353037
Induction of an immune response by oral administration of recombinant botulinum toxin. Nov 1997
- 6/6/60 11986811 PMID: 9268721
Botulinum E toxin light chain does not cleave SNAP-23 and only partially impairs insulin stimulation of GLUT4 translocation in 3T3-L1 cells. Aug 19 1997
- 6/6/61 11956902 PMID: 9236236
Rac1 mediates collapsin-1-induced growth cone collapse. Aug 15 1997
- 6/6/62 11942980 PMID: 9224685
The interaction of synaptic vesicle-associated membrane protein/syntaxobrevin with botulinum neurotoxins D and F. Jun 16 1997
- 6/6/63 11918493 PMID: 9195974
Analysis of a yeast SNARE complex reveals remarkable similarity to the neuronal SNARE complex and a novel function for the C terminus of the SNAP-25 homolog. Sep 3 1997
- 6/6/64 11887878 PMID: 9182707
Sphingosine 1-phosphate stimulates rho-mediated tyrosine phosphorylation of focal adhesion kinase and paxillin in Swiss 3T3 fibroblasts. Jun 1 1997
- 6/6/65 11700899 PMID: 9077452
Involvement of Rho and Rac small G proteins and Rho GDI in Ca²⁺-dependent exocytosis from PC12 cells. Oct 1996
- 6/6/66 11637159 PMID: 8948442
ADP-ribosylation of the GTP-binding protein RhoA blocks cytoplasmic division in human myelomonocytic cells. Jun 15 1995
- 6/6/67 11596817 PMID: 8907197
Activation of *Clostridium botulinum* C3 exoenzyme-catalyzed ADP-ribosylation of RhoA by K⁺ in a Mg²⁺-dependent manner. Jan 1996
- 6/6/68 11451603 PMID: 8702770
Structural determinants of the specificity for synaptic vesicle-associated membrane protein/syntaxobrevin of tetanus and botulinum type B and G neurotoxins. Aug 23 1996
- 6/6/69 11370700 PMID: 8699032
Directing antigen specificity towards botulinum neurotoxin with combinatorial phage display libraries. Jun 21 1996
- 6/6/70 11314299 PMID: 8626670
Evidence that syntaxin 1A is involved in storage in the secretory pathway. May 10 1996
- 6/6/71 11263823 PMID: 8611567
Botulinum neurotoxin C1 cleaves both syntaxin and SNAP-25 in intact and permeabilized chromaffin cells: correlation with its blockade of catecholamine release. Feb 27 1996
- 6/6/72 11255514 PMID: 8557112
The high-affinity binding of *Clostridium botulinum* type B neurotoxin to synaptotagmin II associated with gangliosides GT1b/GD1a. Jan 15 1996
- 6/6/73 11241115 PMID: 8555186
Active site mutation of the C3-like ADP-ribosyltransferase from *Clostridium Inosom-* analysis of glutamic acid 174. Jan 9 1996
- 6/6/74 11175766 PMID: 7479854
Rho protein regulates tight junctions and perijunctional actin organization in polarized epithelia. Nov 7 1995
- 6/6/75 11138155 PMID: 7476433
Preparation of native and recombinant *Clostridium botulinum* C3 ADP-ribosyltransferase and identification of Rho proteins by ADP-ribosylation. 1995
- 6/6/76 11138153 PMID: 7476431
Purification and assay of recombinant C3 transferase. 1995
- 6/6/77 11137711 PMID: 7592744
Synergistic activation of rat brain phospholipase D by ADP-ribosylation factor and rhoA p21, and its inhibition by *Clostridium botulinum* C3 exoenzyme. Oct 27 1995
- 6/6/78 10688027 PMID: 7860636
The t-SNAREs syntaxin 1 and SNAP-25 are present on organelles that participate in synaptic vesicle recycling. Feb 1995
- 6/6/79 10834579 PMID: 7829513
IgA protease from *Neisseria gonorrhoeae* inhibits exocytosis in bovine chromaffin cells like tetanus toxin. Jan 27 1995
- 6/6/80 10753530 PMID: 7525357
Botulinum C3 exoenzyme blocks the tyrosine phosphorylation of p125FAK and paxillin induced by bombesin and endothelin. Nov 14 1994
- 6/6/81 10750183 PMID: 7954816
The small GTP-binding protein Rho regulates a phosphatidylinositol 4-phosphate 5-kinase in mammalian cells. Nov 4 1994
- 6/6/82 10622052 PMID: 8011337
A post-docking role for syntaxobrevin in synaptic vesicle fusion. Jun 1994
- 6/6/83 10600926 PMID: 8198524
ADP-ribosylation of Rho proteins by *Clostridium botulinum* exoenzyme C3 is influenced by phosphorylation of Rho-associated factors. May 15 1994
- 6/6/84 10515628 PMID: 8294407
Proteolysis of SNAP-25 by types E and A botulinum neurotoxins. Jan 21 1994
- 6/6/85 10365611 PMID: 8243676
Botulinum neurotoxins serotypes A and E cleave SNAP-25 at distinct COOH-terminal peptide bonds. Nov 29 1993
- 6/6/86 10142582 PMID: 8385945
Enhancement of *Clostridium botulinum* C3-catalyzed ADP-ribosylation of recombinant rhoA by sodium dodecyl sulfate. Apr 6 1993
- 6/6/87 10109196 PMID: 7680558
Inhibition of PMA-induced, LFA-1-dependent lymphocyte aggregation by ADP-ribosylation of the small molecular weight GTP binding protein. rho. Mar 1993
- 6/6/88 09713853 PMID: 1551445
ADP-ribosylation by *Clostridium botulinum* C3 exoenzyme increases steady-state GTPase activities of recombinant rhoA and rhoB proteins. Feb 3 1992
- 6/6/89 08584865 PMID: 2497029
Identification of rho as a substrate for botulinum toxin C3-catalyzed ADP-ribosylation. Apr 24 1989
- 6/6/90 08464392 PMID: 2492192
The rho gene product expressed in *E. coli* is a substrate of botulinum ADP-ribosyltransferase C3. Jan 16 1989
- 6/5/48 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv. 12/482093 PMID: 9792657
Characterization of the catalytic site of the ADP-ribosyltransferase *Clostridium botulinum* C2 toxin by site-directed mutagenesis.
Barth H, Preiss J C, Hofmann F, Aktories K
Institut für Pharmakologie und Toxikologie der Albert-Ludwigs-Universität Freiburg, D-79104 Freiburg, Germany.
Journal of biological chemistry (UNITED STATES) Nov 6 1998, 273 (45) p29506-11, ISSN 0021-9258 Journal Code: 2985121R Publishing Model Print
Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM
Record type: MEDLINE; Completed Subfile: INDEX MEDICUS
The actin ADP-ribosylating *Clostridium botulinum* C2 toxin is a binary toxin composed of the binding component C2II and the enzyme component C2I. C2I ADP-ribosylates G-actin at arginine 177, resulting in the depolymerization of the actin cytoskeleton. Here, we studied the structure-function relationship of C2I by site-directed mutagenesis. Exchange of Glu389 to glutamine caused the complete loss of ADP-ribosyltransferase and NAD-glycohydrolase activities of C2I. In contrast, exchange of Glu387 to glutamine blocked ADP-ribosyltransferase but not NAD-glycohydrolase activity. Whereas photoaffinity labeling of the double mutant E387Q/E389Q C2I with [carboxyl-¹⁴C]NAD was blocked, labeling of the single C2I mutants was reduced (E389Q) or not changed (E387Q). Exchange of the STS motif (amino acid residues 348-350) of C2I caused a decrease in transferase activity by more than 99 (S348A) and 90% (T349V), or did not affect activity (S350A). Exchange of Arg289 and Arg300 to lysine reduced transferase activity to <0.1 and

approximately 35% of wild-type activity. The data indicate that the amino acid residues Glu389, Glu387, Ser348, and Arg299, which are conserved in various prokaryotic and eukaryotic arginine-modifying ADP-ribosyltransferases, are essential for ADP-ribosyltransferase activity of the enzyme component of *C. botulinum* C2 toxin.

Tags: Research Support, Non-U.S. Gov't

Descriptors: *Botulinum Toxins--metabolism--ME; *Poly(ADP-ribose) Polymerases--metabolism--ME; Amino Acid Sequence; Animals; Botulinum Toxins--chemistry--CH; Botulinum Toxins--genetics--GE; CHO Cells; Catalytic Domain; Hamsters; Humans; Molecular Sequence Data; Mutagenesis, Site-Directed; Photoaffinity Labels; Poly(ADP-ribose) Polymerases--chemistry--CH; Poly(ADP-ribose) Polymerases--genetics--GE; Recombinant Proteins--chemistry--CH; Recombinant Proteins--genetics--GE; Recombinant Proteins--metabolism--ME; Sequence Homology, Amino Acid CAS Registry No.: 0 (Botulinum Toxins); 0 (Photoaffinity Labels); 0 (Recombinant Proteins); 0 (botulinum toxin type C) Enzyme No.: EC 2.4.2.30 (Poly(ADP-ribose) Polymerases) Record Date Created: 1998/12/10 Record Date Completed: 1998/12/10

6/6/02 DIALOG(R)File 155.MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.

12383413 PMID: 9693060

Production and purification of the heavy-chain fragment C of botulinum neurotoxin, serotype B, expressed in the methylotrophic yeast *Pichia pastoris*.

Potter K J; Bevins M A; Vassiliev E V; Chiruvolu V R; Smith T; Smith L A;

Meagher M M

Department of Food Science and Technology, Biological Process Development Facility, University of Nebraska-Lincoln, 68583-0919, USA.

Protein expression and purification (UNITED STATES) Aug 1998, 13 (3) p357-65, ISSN 1046-5928 Journal Code: 9101496 Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM

Record type: MEDLINE; Completed Subfile: INDEX MEDICUS

A recombinant Hc fragment of botulinum neurotoxin, serotype B (rBoNTB(Hc)), has been successfully expressed in a *Mut+* strain of the methylotrophic yeast *Pichia pastoris* for use as an antigen in a proposed human vaccine. The fermentation process consisted of batch phase on glycerol, followed by glycerol and methanol fed batch phases yielding a final cell mass of 60 g/L (dcw) and was easily scaled-up to 60 L. A multistep ion-exchange chromatographic purification process was employed to produce 99% pure Hc fragment. The final yield of the purified antigen was 390 mg per kilogram of wet cell mass. The purified Hc fragment of serotype B was stable, elicited an immune response in mice, and protected upon challenge with native botulin. Copyright 1998 Academic Press.

Tags: Research Support, U.S. Gov't, Non-P.H.S.

Descriptors: *Botulinum Toxins--genetics--GE; *Neurotoxins--genetics--GE; *Pichia--genetics--GE; Amino Acid Sequence; Animals; Blotting, Western; Botulinum Toxins--chemistry--CH; Botulinum Toxins--isolation and purification--IP; Chromatography, Ion Exchange; Cloning, Molecular; Electrophoresis, Polyacrylamide Gel; Mice; Molecular Sequence Data; Neurotoxins--chemistry--CH; Neurotoxins--isolation and purification--IP; Recombinant Proteins--chemistry--CH; Recombinant Proteins--genetics--GE; Recombinant Proteins--isolation and purification--IP CAS Registry No.: 0 (Botulinum Toxins); 0 (Neurotoxins); 0 (Recombinant Proteins) Record Date Created: 1998/09/17 Record Date Completed: 1998/09/17

9/6/1 18520371 PMID: 15933619

Analysis of active site residues of botulinum neurotoxin E by mutational, functional, and structural studies: Glu335Gln is an aspartic residue. Jun 14 2005

9/6/2 18119249 PMID: 15769746

Lipid raft association of SNARE proteins regulates exocytosis in PC12 cells. May 20 2005

9/6/3 15324488 PMID: 14978027

RaA-exocyst interaction mediates GTP-dependent exocytosis. May 7 2004

9/6/4 15103634 PMID: 14663070

Structural analysis by X-ray crystallography and calorimetry of a haemagglutinin component (HA1) of the progenitor toxin from *Clostridium botulinum*. Dec 2003

9/6/5 14920145 PMID: 12911311

Rho-specific *Bacillus cereus* ADP-ribosyltransferase C3 α er cloning and characterization. Aug 19 2003

9/6/6 14845560 PMID: 12796138

Nitric oxide-induced decrease in calcium sensitivity of resistance arteries is attributable to activation of the myosin light chain phosphatase and antagonized by the RhoA/Rho kinase pathway. Jun 24 2003

9/6/7 14772873 PMID: 12731878

Channel formation by the binding component of *Clostridium botulinum* C2 toxin: glutamate 307 of C2II affects channel properties in vitro and pH-dependent C2II translocation in vivo. May 13 2003

9/6/8 14077815 PMID: 11850454

The core membrane fusion complex governs the probability of synaptic vesicle fusion but not transmitter release kinetics. Feb 15 2002

9/6/9 14048603 PMID: 11814347

Structure-function analysis of the Rho-ADP-ribosylating exoenzyme C3stau2 from *Staphylococcus aureus*. Feb 5 2002

9/6/10 14030272 PMID: 11792805

SNAP-25 with mutations in the zero layer supports normal membrane fusion kinetics. Dec 2001

9/6/11 13946422 PMID: 11700044

Site-directed mutagenesis identifies active-site residues of the light chain of botulinum neurotoxin type A. Nov 16 2001

9/6/12 13800583 PMID: 11373287

A discontinuous SNAP-25 C-terminal coil supports exocytosis. Jul 27 2001

9/6/13 13687573 PMID: 11329292

Thermal stabilization of the catalytic domain of botulinum neurotoxin E by phosphorylation of a single tyrosine residue. Feb 20 2001

9/6/14 13520056 PMID: 10488092

Gap14(13) stimulates Rho-dependent activation of the cyclooxygenase-2 promoter. Sep 24 1999

9/6/15 13451445 PMID: 10409113

Protein kinase C inhibits Kv1.1 potassium channel function. Jul 1999

9/6/16 13312586 PMID: 10085240

SNAP-25a and -25b isoforms are both expressed in insulin-secreting cells and can function in insulin secretion. Apr 1 1999

9/6/17 12997956 PMID: 10954418

Membrane localization and biological activity of SNAP-25 cysteine mutants in insulin-secreting cells. Sep 2000

9/6/18 12883746 PMID: 10748216

Recognition of RhoA by *Clostridium botulinum* C3 exoenzyme. Jun 2 2000

9/6/19 12830226 PMID: 10768830

Rho-A is critical for osteoclast podosome organization, motility, and bone resorption. Apr 21 2000

9/6/20 12660268 PMID: 10582602

Botulinum neurotoxin E-insensitive mutants of SNAP-25 fail to bind VAMP but support exocytosis. Dec 1999

9/6/21 12482093 PMID: 9792657

Characterization of the catalytic site of the ADP-ribosyltransferase *Clostridium botulinum* C2 toxin by site-directed mutagenesis. Nov 6 1998

9/6/22 12292473 PMID: 9601102

The small GTP-binding protein Rac-G regulates uric acid formation in the protozoan parasite *Entamoeba histolytica*. Jun 1998

9/6/23 12114293 PMID: 9414082

Botulinum neurotoxin types A and E require the SNARE motif in SNAP-25 for proteolysis. Nov 24 1997

9/6/24 11942980 PMID: 9224685

The interaction of synaptic vesicle-associated membrane protein/synaptobrevin with botulinum neurotoxins D and F. Jun 16 1997

9/6/25 11924409 PMID: 9202074

Rho proteins play a critical role in cell migration during the early phase of mucosal restitution. Jul 1 1997

9/6/26 11680527 PMID: 9003771

Mutational analysis of VAMP domains implicated in Ca²⁺-induced insulin exocytosis. Dec 16 1996

9/6/27 11241115 PMID: 8555186

Active site mutation of the C3-like ADP-ribosyltransferase from *Clostridium imosum*-analysis of glutamic acid 174. Jan 9 1996

9/6/28 11101927 PMID: 7672106

Identification of Glu173 as the critical amino acid residue for the ADP-ribosyltransferase activity of *Clostridium botulinum* C3 exoenzyme. Sep 4 1995

10/6/1 12950160 PMID: 10899856

The C terminus of component C2II of *Clostridium botulinum* C2 toxin is essential for receptor binding. Aug 2000

10/6/2 12469820 PMID: 9782457

On the action of botulinum neurotoxins A and E at cholinergic terminals. Apr 1998

10/6/3 12403522 PMID: 9712688

Ganglioside GT1b as a complementary receptor component for *Clostridium botulinum* neurotoxins. Aug 1998

10/6/4 11179774 PMID: 7578132

Expression and purification of the light chain of botulinum neurotoxin A: a single mutation abolishes its cleavage of SNAP-25 and neurotoxicity after reconstitution with the heavy chain. Nov 21 1995

10/6/5 10163183 PMID: 8484722

ADP-ribosylation of *Drosophila* indirect-flight-muscle actin and arthrin by *Clostridium botulinum* C2 toxin and *Clostridium perfringens* iota toxin. Apr 15 1993

11/6/1 15332747 PMID: 15133114

Molecular characterization of binding subcomponents of *Clostridium botulinum* type C progenitor toxin for intestinal epithelial cells and erythrocytes. May 2004

11/6/2 15070121 PMID: 14622125

Lysophosphatidic acid induces process retraction in CG-4 line oligodendrocytes and oligodendrocyte precursor cells but not in differentiated oligodendrocytes. Nov 2003

11/6/3 14451966 PMID: 12391898

[Botulinism: the agent, mode of action of the botulinum neurotoxins, forms of acquisition, treatment and prevention] Le botulisme: agent, mode d'action des neurotoxines botuliques, formes d'acquisition, traitement et prevention. Aug 2002

11/6/4 13812149 PMID: 11478959

Botulinum neurotoxin types B and E: purification, limited proteolysis by endoprotease Glu-C and pepsin, and comparison of their identified cleaved sites relative to the three-dimensional structure of type A neurotoxin. Oct 2001

11/6/5 13765559 PMID: 11427819

[Mechanism of action and therapeutic uses of botulinum and tetanus neurotoxins] Mechanismes d'action et utilisations thérapeutiques des neurotoxines botuliques et tétaniques. May 2001

11/6/6 12979051 PMID: 10937461

The role of drug treatment in children with strabismus and amblyopia. Mar-Apr 2000

11/6/7 12404747 PMID: 9717738

Covalent structure of botulinum neurotoxin type B: location of sulfhydryl groups and disulfide bridge and identification of C-termini of light and heavy chains. Jul 1998

11/6/8 10357199 PMID: 7901925

- Tetanus and botulinum neurotoxins: a new group of zinc proteases. Sep 1993
- 11/6/9 08318311 PMID: 3398838
Circular dichroic and fluorescence spectroscopic study of the conformation of botulinum neurotoxin types A and E. Feb 1988
- 11/6/10 07740340 PMID: 3538716
Molecular characterization of a protein, insoluble at low temperature, produced by Clostridium botulinum type G. Aug 1986
- 11/6/11 07042616 PMID: 6382680
Purification and amino acid composition of type A botulinum neurotoxin. 1984
- 11/6/12 06327300 PMID: 7080765
[Characteristics of the activation of type E, Cl botulinum toxins] Nekotorye osobennosti aktivatsii botulinicheskogo toksina tipa E. Apr 1982
- 11/6/13 05629302 PMID: 115504
[Isolation and properties of highly purified C1 botulinum toxin type E] Vydelenie i nekotorye svoistva vysokochisticheskogo botulinicheskogo toksina tipa E. Aug 1979
- 11/7/14 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv. 13812149 PMID: 11478959
Botulinum neurotoxin types B and E: purification, limited proteolysis by endoproteinase Glu-C and pepsin, and comparison of their identified cleaved sites relative to the three-dimensional structure of type A neurotoxin.
Prabakaran S, Tepp W, DasGupta B R
Department of Food Microbiology and Toxicology, University of Wisconsin, Madison, 53706, USA.
- Toxicon - official journal of the International Society on Toxinology (England) Oct 2001, 39 (10) p1515-31, ISSN 0041-0101 Journal Code: 1307333
Contract/Grant No.: NS 17742; NS; NINDS Publishing Model Print
Document type: MEDLINE; Completed
Record type: MEDLINE; Completed
Botulinum neurotoxin (NT) serotypes B and E are approximately 150 kDa proteins. Isolated from the liquid culture of Clostridium botulinum the NT type E is a single chain protein while the NT type B, from the proteolytic strain of the bacteria, is a mixture of dichain (nicked within a disulfide loop located about one-third the way from the N-terminus to the C-terminus) protein and its precursor single-chain protein. Endoproteinase Glu-C (EC 3.4.21.19) and pepsin (EC 3.4.23.1) were used for controlled digestion of NT types B and E; the amino acid residues flanking many of the cleavable peptide bonds were identified and the corresponding proteolytic fragments partially characterized. Chemical identification of 82 and 108 residues of types B and E NT, respectively, revealed that the residue 738 and 1098 in type E NT, identified as Leu and Asn, respectively, differ from those deduced from nucleotide sequences. Several fragments overlapped spanning various segments of the NT's functional domains; they appear to have potential for structure-function studies of the NT. The cleavage sites were compared with the previously determined proteolyzed sites on NT types A and E. The cleavage sites of the NT types A, B and E, all exposed on the protein surface, were scrutinized in the context of the three-dimensional structure of crystallized NT type A [Lacy, D.B., Stevens, R.C., 1999. J. Mol. Biol. 291, 1091-1104]. Detailed procedures for isolation of pure NT types B and E in large quantities (average yield 92 and 62 mg, respectively) suitable for crystallization are reported.
Record Date Created: 20010731 Record Date Completed: 20011004
- 11/5/8 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv. 10357199 PMID: 7901925
Tetanus and botulinum neurotoxins: a new group of zinc proteases.
Montecucco C, Schiavo G
Department of Biomedical Sciences, University of Padova, Italy.
Trends in biochemical sciences (ENGLAND) Sep 1993, 18 (9) p324-7, ISSN 0968-0004 Journal Code: 7610674
- Publishing Model Print Document type: Journal Article; Review; Tutorial
Languages: ENGLISH
Main Citation Owner: NLM Record type: MEDLINE; Completed Subfile: INDEX
MEDICUS
The active forms of tetanus and botulinum neurotoxins, released from the precursor molecule by specific proteolysis and reduction, block the release of neurotransmitters via a Zn(2+)-dependent protease activity. VAMP/synaptobrevin, an integral membrane protein of the synaptic vesicles, is cleaved at a single site by tetanus and botulinum B, D and F neurotoxins. The unique sequence, mechanism of activation and site of activity of clostridial neurotoxins mark them out as an independent group of Zn(2+)-endopeptidases. (38 Refs.)
Tags: Research Support, Non-U.S. Gov't
CAS Registry No.: 0 (Botulinum Toxins); 0 (Membrane Proteins); 0 (Nerve Tissue Proteins); 0 (Neurotransmitters); 0 (Tetanus Toxin); 0 (vesicle-associated membrane protein)
Enzyme No.: EC 3.4.24 (Metalloendopeptidases); EC 3.4.24. (zinc-endopeptidase, tetanus neurotoxin)
Record Date Created: 19931214 Record Date Completed: 19931214
- 11/7/10 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv. 07740340 PMID: 3538716
Molecular characterization of a protein, insoluble at low temperature, produced by Clostridium botulinum type G.
Gimenez JA, Cascone O, Biscoglio de Jimenez Bonino M J, Paladini A C
Zentralblatt für Bakteriologie, Mikrobiologie, und Hygiene. Series A, Medical microbiology, infectious diseases, virology, parasitology (GERMANY, WEST) Aug 1986, 262 (2) p179-88, ISSN 0176-6724 Journal Code: 8403032
Publishing Model Print Document type: Journal Article Languages: ENGLISH
Main Citation Owner: NLM
Record type: MEDLINE; Completed
A preliminary study of a low-toxicity protein, called cryoprotein, produced by Clostridium botulinum type G, led to a better characterization of this substance and to discriminate its relationship with type G botulinum toxin. This sparingly soluble protein has been characterized as an aggregated form of a soluble precursor with an Mr of 170 000. This phenomenon is temperature-dependent. The monomeric protein is usually contaminated with a lower Mr form (150 000) quite probably originated by a limited proteolytic process. The amino acid composition of this protein is relatively analogous to that of the botulinum toxins A and B, the only notable exception being the absence of cysteine. The N-terminal amino acid is alanine and the C-terminal sequence is Val-Ala-Leu-OH. The low toxicity which is usually demonstrable in samples of this protein disappears after a reductive treatment, strongly suggesting that it is not an intrinsic property. Taking into account that some of its physicochemical properties are similar to those of the known botulinum toxins, it is quite probable that this substance accompanies G toxin preparations currently obtained by routine methods, increasing its non-toxic antigenic mass. This fact could be critical to the sensitivity and specificity of G toxin immunological detection methods.
Record Date Created: 19870102 Record Date Completed: 19870102
- 11/7/11 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv. 07042616 PMID: 6382680
Purification and amino acid composition of type A botulinum neurotoxin.
DasGupta B R, Sathiyamoorthy V
Toxicon - official journal of the International Society on Toxinology (ENGLAND) 1984, 22 (3) p415-24, ISSN 0041-0101 Journal Code: 1307333 Contract/Grant No.: NS17742; NS; NINDS Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM Record type: MEDLINE; Completed
A method to purify type A botulinum neurotoxin from a 64 liter bacterial culture is reported. The procedure includes cation exchange chromatography at pH 7.0. The final product, essentially homogeneous (according to polyacrylamide gel-sodium dodecylsulfate electrophoresis), is a mixture of two forms of the neurotoxin (mol. wt 145,000), the dichain or nicked form (over 95%) and its precursor the single chain or
- unnicked form. Two batches of the neurotoxin purified by the method described here and one batch purified according to the method of Sugii and Sakaguchi were similar in purity and amino acid composition. The best estimate of the number of amino acid residues per neurotoxin molecule (mol. wt 145,000) is:
Asp200Thr75Ser79Glu114Pro44Gly64Ala53Val70Cys10Met22Ile111Leu104Tyr71Phe68 Lys100His14Arg43Trp17.
Record Date Created: 19841012 Record Date Completed: 19841012
- 13/6/1 17945879 PMID: 15683238
A spider toxin that induces a typical effect of scorpion alpha-toxins but competes with beta-toxins on binding to insect sodium channels. Feb 8 2005
- 13/6/2 17391913 PMID: 15147240
Topology of the substrate-binding site of a Lys49-phospholipase A2 influences Ca2+-independent membrane-damaging activity. Aug 15 2004
- 13/6/3 16325557 PMID: 15488774
Basic amino acid residues in the beta-structure region contribute, but not critically, to presynaptic neurotoxicity of ammotoxin A. Nov 1 2004
- 13/6/4 15272414 PMID: 15054090
Conversion of a scorpion toxin agonist into an antagonist highlights an acidic residue involved in voltage sensor trapping during activation of neuronal Na+ channels. Apr 2004
- 13/6/5 15222801 PMID: 14992587
Resiniferatoxin binds to the capsaicin receptor (TRPV1) near the extracellular side of the S4 transmembrane domain. Mar 9 2004
- 13/6/6 15210197 PMID: 14672947
Dissection of the functional surface of an anti-insect excitatory toxin illuminates a putative 'hot spot' common to all scorpion beta-toxins affecting Na+ channels. Feb 27 2004
- 13/6/7 14751905 PMID: 12705833
Exporation of the functional site of a scorpion alpha-like toxin by site-directed mutagenesis. Apr 29 2003
- 13/6/8 14666413 PMID: 12595696
Structural analysis of Tityus serrulatus Ts1 neurotoxin at atomic resolution: insights into interactions with Na+ channels. Mar 2003
- 13/6/9 14445108 PMID: 12383115
Roles of disulfide bridges in scorpion toxin BmK M1 analyzed by mutagenesis. Nov 2002
- 13/6/10 14148995 PMID: 11931685
Phenylalanine-24 in the N-terminal region of ammodytoxin is important for both enzymic activity and presynaptic toxicity. Apr 15 2002
- 13/6/11 14138559 PMID: 11920913
The active site of human paraoxonase (PON1). Dec 2001
- 13/6/12 13949849 PMID: 11527975
Design and characterization of a highly selective peptide inhibitor of the small conductance calcium-activated K+ channel, SKCa2. Nov 16 2001
- 13/6/13 13705739 PMID: 11352729
Interaction of a toxin from the scorpion Tityus serrulatus with a cloned K+ channel from squid (sqKv1A). May 22 2001
- 13/6/14 13416021 PMID: 10377255
An aromatic, but not a basic, residue is involved in the toxicity of group-II phospholipase A2 neurotoxins. Jul 1 1999
- 13/6/15 13319140 PMID: 10092644
Subunit interface selectivity of the alpha-neurotoxins for the nicotinic acetylcholine receptor. Apr 2 1999
- 13/6/16 13319087 PMID: 10092591
Importance of the carboxyl terminus in the folding and function of alpha-hemolysin of Staphylococcus aureus. Apr 2 1999

- 13/6/17 13028016 PMID: 10987857
Structural implications on the interaction of scorpion alpha-like toxins with the sodium channel receptor site inferred from toxin iodination and pH-dependent binding. Oct 2000
- 13/6/18 12967939 PMID: 10807914
Relative spatial position of a snake neurotoxin and the reduced disulfide bond alpha (Cys192-Cys193) at the alpha gamma interface of the nicotinic acetylcholine receptor. Aug 18 2000
- 13/6/19 12969238 PMID: 10920011
Mechanisms of neurotoxin action on Shaker potassium channels. Aug 2000
- 13/6/20 12873054 PMID: 10814779
Modulation of Alzheimer's beta-amyloid neurotoxicity by site-directed single-chain antibody. Jul 1 2000
- 13/6/21 12479802 PMID: 9792175
Effect of site directed mutagenesis on the activity of recombinant trimucrotoxin, a neurotoxic phospholipase from *Trimeresurus mucrosquamatus* venom. Nov 1998
- 13/6/22 12467470 PMID: 9778362
A snake toxin inhibitor of inward rectifier potassium channel ROMK1. Oct 20 1998
- 13/6/23 12048635 PMID: 9341237
Structure and function of *Cerebratulus lacteus* neurotoxin B-IV: tryptophan-30 is critical for function while lysines-18, -19, -29, and -33 are not required. Oct 28 1997
- 13/6/24 12042041 PMID: 9335541
Nonidentity of the alpha-neurotoxin binding sites on the nicotinic acetylcholine receptor revealed by modification in alpha-neurotoxin and receptor structures. Oct 21 1997
- 13/6/25 11901535 PMID: 9179299
In vitro folding and functional analysis of an anti-insect selective scorpion depressant neurotoxin produced in *Escherichia coli*. Jun 1997
- 13/6/26 11892865 PMID: 9169449
Identification of structural elements of a scorpion alpha-neurotoxin important for receptor site recognition. Jun 6 1997
- 13/6/27 11818631 PMID: 9116014
Critical interactions at the dimer interface of kappa-bungarotoxin, a neuronal nicotinic acetylcholine receptor antagonist. Mar 18 1997
- 13/6/28 11801672 PMID: 9054546
Solution structures of a highly insecticidal recombinant scorpion alpha-toxin and a mutant with increased activity. Mar 4 1997
- 13/6/29 11690643 PMID: 9027980
Binding of native kappa-neurotoxins and site-directed mutants to nicotinic acetylcholine receptors. Nov-Dec 1996
- 13/6/30 11627756 PMID: 8939911
Role of electrostatic interactions in defining the potency of neurotoxin B-IV from *Cerebratulus lacteus*. Nov 22 1996
- 13/6/31 11607874 PMID: 8916901
Role for Pro-13 in directing high-affinity binding of anthopleurin B to the voltage-sensitive sodium channel. Nov 12 1996
- 13/6/32 11419633 PMID: 8756487
Functional expression and genetic alteration of an alpha scorpion neurotoxin. Aug 6 1996
- 13/6/33 11398988 PMID: 8731479
Glutamate receptor ion channel properties predict vulnerability to cytotoxicity in a transfected nonneuronal cell line. Feb 1996
- 13/6/34 11342972 PMID: 8639500
Importance of highly conserved anionic residues and electrostatic interactions in the activity and structure of the cardiotonic polypeptideanthopleurin B. Mar 19 1996
- 13/6/35 10940534 PMID: 7721859
Genetic engineering of snake toxins. The functional site of Erabutoxin a, as delineated by site-directed mutagenesis, includes variant residues. Apr 21 1995
- 13/6/36 10809137 PMID: 7605827
On the site by which alpha-dendrotoxin binds to voltage-dependent potassium channels: site-directed mutagenesis reveals that the lysine triplet 28-30 is not essential for binding. Dec 19 1994
- 13/6/37 10607472 PMID: 7911329
A single mutation in the recombinant light chain of tetanus toxin abolishes its proteolytic activity and removes the toxicity seen after reconstitution with native heavy chain. Jun 7 1994
- 13/6/38 10548224 PMID: 8142390
Site-directed mutagenesis of kappa-bungarotoxin: implications for neuronal receptor specificity. Apr 5 1994
- 13/6/39 10513454 PMID: 8288644
Role of the cationic residues arginine 14 and lysine 48 in the function of the cardiotonic polypeptide anthopleurin B. Jan 14 1994
- 13/6/40 10282717 PMID: 8103516
Role of the large extracellular domain of metabotropic glutamate receptors in agonist selectivity determination. Sep 15 1993
- 13/6/41 10110080 PMID: 7680721
Receptor sites for open channel blockers of Shaker voltage-gated potassium channels--molecular approaches. 1993
- 13/6/42 09943580 PMID: 1409549
Insertion of a disulfide-containing neurotoxin into *E. coli* alkaline phosphatase: the hybrid retains both biological activities. Apr 1992
- 13/6/43 09887742 PMID: 1511681
Role and environment of the conserved Lys27 of snake curaremimetic toxins as probed by chemical modifications, site-directed mutagenesis and photolabelling experiments. Aug 15 1992
- 13/6/44 09745101 PMID: 1568473
The three-dimensional structure of norexin, a presynaptic neurotoxic phospholipase A2 at 2.0 Å resolution. Apr 20 1992
- 13/6/45 09453747 PMID: 2071577
Mutagenesis of *Cerebratulus lacteus* neurotoxin B-IV identifies NH2-terminal sequences important for biological activity. Jul 15 1991
- 14/6/1 18044823 PMID: 15941066
Molecular cloning and nucleotide sequence analysis of encoded anti-insect toxin BotIT2 from the scorpion *Buthus occitanus tunetanus* venom. 2003
- 14/6/2 17822062 PMID: 15548530
Jingzhaoxin-I, a novel spider neurotoxin preferentially inhibiting cardiac sodium channel inactivation. Apr 1 2005
- 14/6/3 17648343 PMID: 15698615
Endogenous Nix2-HOlg2+ oligodendrocyte precursor cells fail to remyelinate the demyelinated adult rat spinal cord in the absence of astrocytes. Mar 2005
- 14/6/4 17638671 PMID: 15730236
Comparative analysis of a neurotoxin from *Callisotoma canaliculatum* by on-line capillary isotachopheresis/¹H NMR and diffusion ¹H NMR. Feb 2005
- 14/6/5 17536836 PMID: 15670783
An unusual soluble beta-tum-rich conformation of prion is involved in fibril formation and toxic to neuronal cells. Mar 4 2005
- 14/6/6 16400836 PMID: 15217615
Structure and biosynthesis of the jamaicamides, new mixed polyketide-peptide neurotoxins from the marine cyanobacterium *Lyngbya majuscula*. Jun 2004
- 14/6/7 15487588 PMID: 15388848
Efficient reversal of Alzheimer's disease fibril formation and elimination of neurotoxicity by a small molecule. Oct 5 2004
- 14/6/8 15381346 PMID: 15197738
Characterization of V642I-AbetaPP-induced cytotoxicity in primary neurons. Jul 1 2004
- 14/6/9 15379119 PMID: 15084603
Jingzhaoxin-III, a novel spider toxin inhibiting activation of voltage-gated sodium channel in rat cardiac myocytes. Jun 18 2004
- 14/6/10 15352700 PMID: 15158020
Protective effects of S-nitrosoglutathione against neurotoxicity of 3-nitropropionic acid in rat. May 27 2004
- 14/6/11 15315383 PMID: 15109888
Isolation, molecular cloning and functional characterization of a novel beta-toxin from the Venezuelan scorpion, *Tityus zuluensis*. May 2004
- 14/6/12 15306673 PMID: 15095690
The presence of the APP(swe) mutation in mice does not increase the vulnerability of cholinergic basal forebrain neurons to neuroinflammation. 2004
- 14/6/13 15183557 PMID: 14757201
cDNA sequence analysis of seven peptide toxins from the spider *Selenocosmia huwena*. Dec 2003
- 14/6/14 15149327 PMID: 14716025
Interactive effects of excitotoxic injury and dietary restriction on microglia and neurogenesis in the hippocampus of adult mice. 2003
- 14/6/15 14973428 PMID: 13130517
Overexpression of amyloid precursor protein is associated with degeneration, decreased viability, and increased damage caused by neurotoxins (prostaglandins A1 and E2, hydrogen peroxide, and nitric oxide) in differentiated neuroblastoma cells. Oct 1 2003
- 14/6/16 14970167 PMID: 12972326
Isolation of scorpion (*Androctonus amoreuxi*) putative alpha neurotoxins and parallel cloning of their respective cDNAs from a single sample of venom. Sep 15 2003
- 14/6/17 14907038 PMID: 12893066
Implication of brevetoxin B1 and P6Tx-3 in neurotoxic shellfish poisoning in New Zealand by isolation and quantitative determination with liquid chromatography-tandem mass spectrometry. Jul 2003
- 14/6/18 14843167 PMID: 12818191
A novel class of peptide found in scorpion venom with neurodepressant effects in peripheral and central nervous system of the rat. Jun 26 2003
- 14/6/19 14810268 PMID: 12777702
Endogenous estrogen formation is neuroprotective in model of cerebellar ataxia. Jun 2003
- 14/6/20 147556336 PMID: 12711370
Developmental expression of the cell cycle and apoptosis controlling gene, *Lot1*, in the rat cerebellum and in cultures of cerebellar granule cells. May 14 2003
- 14/6/21 14685473 PMID: 12621621
Liquid chromatography with electrospray ion-trap mass spectrometry for the determination of anatoxins in cyanobacteria and drinking water. 2003
- 14/6/22 14643321 PMID: 12560130
Serotonin depletion in rat hippocampus attenuates L-NAME-induced spatial learning deficits. Feb 14 2003
- 14/6/23 14567925 PMID: 12566925
Molecular mechanisms for neuronal cell death by Alzheimer's amyloid precursor protein-relevant insults. Sep-Oct 2002
- 14/6/24 14546635 PMID: 12505425
In vitro studies of Flemish, Dutch, and wild-type beta-amyloid provide evidence for two-staged neurotoxicity. Nov 2002
- 14/6/25 14445826 PMID: 12383342

- The beta-amyloid precursor protein (APP) and Alzheimer's disease: does the tail wag the dog? Nov 2002
- 14/6/26 14343795 PMID: 12168693
Characterization and gene organization of Taiwan banded krait (*Bungarus multicinctus*) gamma-bungarotoxin. May 2002
- 14/6/27 14314829 PMID: 12128077
Correlation between beta-amyloid peptide production and human APP-induced neuronal death. Jul 2002
- 14/6/28 14310708 PMID: 12123858
Ergothioneine treatment protects neurons against N-methyl-D-aspartate excitotoxicity in an in vivo rat retinal model. Aug 2 2002
- 14/6/29 14199154 PMID: 11997884
Distinct subsets of nucleus basalis neurons exhibit similar sensitivity to excitotoxicity. May 7 2002
- 14/6/30 14186990 PMID: 11982636
Neural cells from primary human striatal xenografts migrate extensively in the adult rat CNS. Apr 2002
- 14/6/31 14183581 PMID: 11861655
Intracellular amyloid-beta 1-42, but not extracellular soluble amyloid-beta peptides, induces neuronal apoptosis. May 3 2002
- 14/6/32 14125500 PMID: 11905990
Neurotrophic mechanisms triggered by Alzheimer's disease-linked mutant M146L presenilin 1: involvement of NO synthase via a novel pertussis toxin target. Feb 2002
- 14/6/33 14087201 PMID: 11860470
Glia-derived arginine, the nitric oxide precursor, protects neurons from NMDA-induced excitotoxicity. Dec 2001
- 14/6/34 13983820 PMID: 11735408
Slow folding of three-fingered toxins is associated with the accumulation of native disulfide-bonded intermediates. Dec 18 2001
- 14/6/35 13944368 PMID: 11698110
Molecular cloning and characterization of four scorpion K(+) toxin-like peptides: a new subfamily of venom peptides (alpha-KTx14) and genomic analysis of a member. Sep 2001
- 14/6/36 13932320 PMID: 11684050
Neurotoxic and neuroprotective effects of glutamate are enhanced by introduction of amyloid precursor protein cDNA. Nov 9 2001
- 14/6/37 13909092 PMID: 11597611
Kainate excitotoxicity in organotypic hippocampal slice cultures: evidence for multiple apoptotic pathways. Oct 19 2001
- 14/6/38 13892608 PMID: 11578751
The molecular bases of Alzheimer's disease and other neurodegenerative disorders. Sep-Oct 2001
- 14/6/39 13869615 PMID: 11553289
NAIP protects the nigrostriatal dopamine pathway in an intrastriatal 6-OHDA rat model of Parkinson's disease. Jul 2001
- 14/6/40 13862414 PMID: 11544042
The use of GABA(A) receptors expressed in neural precursor cells for cell-based assays. Sep 2001
- 14/6/41 13860256 PMID: 11531526
Cloning and characterization of the pseudonajatoxin b precursor. Sep 15 2001
- 14/6/42 13812156 PMID: 11478966
Characterisation of the gene encoding the alpha-toxin Amm V from the scorpion *Androctonus maurelaticus maurelaticus*. Oct 2001
- 14/6/43 13796246 PMID: 11461960
Homocysteine potentiates beta-amyloid neurotoxicity: role of oxidative stress. Jul 2001
- 14/6/44 13770996 PMID: 11432978
C-terminal fragment of amyloid precursor protein induces astrocytosis. Jul 2001
- 14/6/45 13704241 PMID: 11351342
Brain aromatase is neuroprotective. Jun 15 2001
- 14/6/46 13661506 PMID: 11298767
Molecular characterization of an anti-epilepsy peptide from the scorpion *Buthus martensii* Karsch. Apr 2001
- 14/6/47 13616181 PMID: 11238735
Homocysteine potentiates copper- and amyloid beta peptide-mediated toxicity in primary neuronal cultures: possible risk factors in the Alzheimer's-type neurodegenerative pathways. Mar 2001
- 14/6/48 13594342 PMID: 11158256
Activation of microglia by secreted amyloid precursor protein evokes release of glutamate by cystine exchange and attenuates synaptic function. Feb 2001
- 14/6/49 13588012 PMID: 11172780
White matter damage following systemic injection of the mitochondrial inhibitor 3-nitropropionic acid in rat. Feb 23 2001
- 14/6/50 13567305 PMID: 11125240
Neurotogenic-neurotoxic effects of membrane-associated forms of amyloid precursor protein. Jan-Feb 2001
- 14/6/51 13559996 PMID: 10978740
Molecular cloning and sequence analysis of cDNAs encoding a beta-toxin-like peptide and two MKTx I homologues from scorpion *Buthus martensii* Karsch. Feb-Mar 2001
- 14/6/52 13475030 PMID: 10443484
Enzymatic formation of 2,3-diaminopropionic acid, the direct precursor of the neurotoxin beta-ODAP, in *Lathyrus sativus*. Jul 1999
- 14/6/53 13460192 PMID: 10428048
Secreted form of amyloid precursor protein enhances basal glucose and glutamate transport and protects against oxidative impairment of glucose and glutamate transport in synaptosomes by a cyclic GMP-mediated mechanism. Aug 1999
- 14/6/54 13433427 PMID: 10400300
Molecular characterization of a new excitatory insect neurotoxin with an analgesic effect on mice from the scorpion *Buthus martensii* Karsch. Aug 1999
- 14/6/55 13417732 PMID: 10380958
Cation selective channels formed by a C-terminal fragment of beta-amyloid precursor protein. May 14 1999
- 14/6/56 13338185 PMID: 10202534
Etiology and pathogenesis of Parkinson's disease. 1999
- 14/6/57 13304073 PMID: 10078863
The cDNA sequence of an excitatory insect selective neurotoxin from the scorpion *Buthus martensii* Karsch. Feb 1999
- 14/6/58 13282741 PMID: 10024130
Muscle precursor cells injected into irradiated mdx mouse muscle persist after serial injury. Feb 1999
- 14/6/59 13139354 PMID: 11124892
Antibody dy-regulated neurotoxic function of cell-surface beta-amyloid precursor protein. Dec 2000
- 14/6/60 13106698 PMID: 11083227
The lipid peroxidation by-product 4-hydroxynonenal is toxic to axons and oligodendrocytes. Nov 2000
- 14/6/61 13095525 PMID: 11068140
Increased amyloid precursor protein expression and serotonergic sprouting following excitotoxic lesion of the rat magnocellular nucleus basalis: neuroprotection by Ca(2+) antagonist nimodipine. 2000
- 14/6/62 13087240 PMID: 11059270
[Expression and characterization of two kinds of recombinant snake neurotoxins] May 2000
- 14/6/63 13004169 PMID: 10960598
Neurotoxic APP C-terminal and beta-amyloid domains colocalize in the nuclei of substantia nigra pars reticulata neurons undergoing delayed degeneration. Aug 26 2000
- 14/6/64 12942178 PMID: 10894172
Dexamethasone-induced stimulation of glutamine oxidation in astrocytes: a possible mechanism for its neuroprotective activity. Jun 2000
- 14/6/65 12901650 PMID: 10850733
Interaction of amyloid precursor proteins and heme oxygenase. 2000
- 14/6/66 12794803 PMID: 10728828
Cloning and characterization of the cDNA sequences of two venom peptides from Chinese scorpion *Buthus martensii* Karsch (Bmk). Jul 2000
- 14/6/67 12780322 PMID: 10708793
Cloning and characterization of a cDNA sequence encoding the precursor of a chbotoxin-like peptide from the Chinese scorpion *Buthus martensii* Karsch. Aug 2000
- 14/6/68 12689517 PMID: 10613517
Effects of in vivo sodium azide administration on the immunohistochemical localization of kynurenine aminotransferase in the rat brain. 1999
- 14/6/69 12648362 PMID: 10569632
Local striatal infusion of MPP+ does not result in increased hydroxylation after systemic administration of 4-hydroxybenzoate. Nov 1999
- 14/6/70 12645931 PMID: 10564538
Dehydroepiandrosterone protects hippocampal neurons against neurotoxin-induced cell death: mechanism of action. Nov 1999
- 14/6/71 12642865 PMID: 10563207
[Cloning and structure of gene encoded alpha-latrotoxin from the Black widow spider venom] [Klorovani i struktura gena, kodirajushchego alpha-latrokristoloksin v sostave iadovitjkh zhelez pauka karakura. Jul 1999
- 14/6/72 12595930 PMID: 10079602
HIV-1-infection in the CNS: A pathogenesis of some neurological syndromes in the light of recent investigations. 1998
- 14/6/73 12524242 PMID: 9839668
Cloning of cDNAs encoding neurotoxic peptides from the spider *Phoneutria nigriventer*. Dec 1998
- 14/6/74 12512694 PMID: 9825941
Caspases dependent DNA fragmentation might be associated with excitotoxicity in Alzheimer disease. Nov 1998
- 14/6/75 12478034 PMID: 9790168
Prostaglandins as putative neurotoxins in Alzheimer's disease. Nov 1998
- 14/6/76 12452357 PMID: 9763469
Induction of NF-kappaB activity during haloperidol-induced oxidative toxicity in cbnol hippocampal cells: suppression of NF-kappaB and neuroprotection by antioxidants. Oct 15 1998
- 14/6/77 12408485 PMID: 9720974
The effects of excitotoxicity on the expression of the amyloid precursor protein gene in the brain and its modulation by neuroprotective agents. 1998
- 14/6/78 12288627 PMID: 9600201
A4 amyloid is toxic to neuronal processes. Jan 1998
- 14/6/79 12282615 PMID: 9593594
Proconvulsant effects of neurosteroids pregnenolone sulfate and dehydroepiandrosterone sulfate in mice. Mar 12 1998
- 14/6/80 12262798 PMID: 9572056

| | | | |
|--|--|--|--|
| Molecular pathogenesis of sporadic and familial forms of Alzheimer's disease. Apr 1998 | | The serpin-enzyme complex receptor recognizes soluble, nontoxic amyloid-beta peptide but not aggregated, cytotoxic amyloid-beta peptide. Jul 26 1996 | 14/6/120 10047818 PMID: 8421518 Neurotoxicity of beta-amyloid. Jan 14 1993 |
| 14/6/81 12256084 PMID: 9566620 | | 14/6/100 11289963 PMID: 8624101 beta-APP cognitive function versus beta-amyloid--induced cell death. Jan 17 1996 | 14/6/121 10006660 PMID: 1458313 Accumulation of amyloid precursor protein in damaged neuronal processes and microglia following intracerebral administration of aluminum salts. Oct 9 1992 |
| 14/6/82 12208357 PMID: 9515959 | | 14/6/101 11262241 PMID: 8580732 The cellular events of injured muscle regeneration depend on the nature of the injury. Nov 1995 | 14/6/122 09852468 PMID: 1637045 The etiology of Alzheimer's disease: the pathogenesis of dementia. The role of neurotoxins. May 11 1992 |
| 14/6/83 12150533 PMID: 9453568 | | 14/6/102 11170853 PMID: 7488136 17-beta estradiol protects neurons from oxidative stress-induced cell death in vitro. Nov 13 1995 | 14/6/123 09776569 PMID: 1814634 Biosynthesis of beta-(isoxazolin-5-on-2-yl)alanine, the precursor of the neurotoxic amino acid beta-N-oxalyL-alpha,beta-diaminopropionic acid. Dec 1991 |
| Endogenously occurring beta-carbofene induces parkinsonism in nonprimate animals: a possible causative protein in idiopathic Parkinson's disease. Feb 1998 | | 14/6/103 11072489 PMID: 7645452 Amyloid precursor protein processing is stimulated by metabotropic glutamate receptors. Aug 15 1995 | 14/6/124 09694018 PMID: 1796479 The cDNA sequence of a depressant insect selective neurotoxin from the scorpion <i>Buthotus judacis</i> . 1991 |
| 14/6/84 11962630 PMID: 9266422 | | 14/6/104 11011639 PMID: 7790874 Apoptotic cell death induced by beta-amyloid 1-42 peptide is cell type dependent. Jul 1995 | 14/6/125 09634920 PMID: 1731777 Overexpression of amyloid precursor protein alters its normal processing and is associated with neurotoxicity. Jan 15 1992 |
| 14/6/85 11963991 PMID: 9248001 | | 14/6/105 10977438 PMID: 7756304 Promoter structure and intron-exon organization of a scorpion alpha-toxin gene. May 23 1995 | 14/6/126 09515824 PMID: 1832593 Dopamine D1 receptor behavioral responsiveness following selective lesions of the striatal patch compartment during development. Jun 21 1991 |
| The CDNA and genomic DNA sequences of a mammalian neurotoxin from the scorpion <i>Buthus martensii</i> Karsch. Jul 1997 | | 14/6/106 10855888 PMID: 7849528 Amyloid beta-protein precursor : new clues to the genesis of Alzheimer's disease. Oct 1994 | 14/6/127 09513256 PMID: 1909818 Calcium mediation of cyanide-induced catecholamine release: implications for neurotoxicity. Sep 1 1991 |
| 14/6/87 11934504 PMID: 9215699 | | 14/6/107 10767109 PMID: 7970221 Enhanced cytotoxicity of amyloid beta-peptide by a complement dependent mechanism. Jul 4 1994 | 14/6/128 9369219 PMID: 1826481 Striatal D1 dopamine receptor morphochemistry following continuous or intermittent L-dopa replacement therapy. Apr 1991 |
| P-P and beta-amyloid fragments activate different neurotoxic mechanisms in cultured mouse cells. Jun 1997 | | 14/6/108 10719181 PMID: 7925845 Induction of beta-amyloid-containing polypeptides in hippocampus: evidence for a concomitant loss of synaptic proteins and interactions with an excitotoxin. Sep 1994 | 14/6/129 09288775 PMID: 2101746 Sensitive murine model and putative antibodies for behavioral toxicosis from contaminated mussel extracts. Sep 1990 |
| 14/6/88 11922831 PMID: 9202294 | | 14/6/109 10869461 PMID: 8055934 Production of active, insect-specific scorpion neurotoxin in yeast. Jul 15 1994 | 14/6/130 09180810 PMID: 2249684 Structure of the snake short-chain neurotoxin, erabutoxin c, precursor gene. Nov 13 1990 |
| 14/6/89 11902026 PMID: 9179790 | | 14/6/110 10535630 PMID: 7510258 Isolation and characterization of two genes encoding caltioxins, neurotoxic peptides from <i>Callitaxis parasitica</i> (Oridaria). Jan 28 1994 | 14/6/131 09084413 PMID: 2387406 Characterization of the transcript for a depressant insect selective neurotoxin gene with an isolated cDNA clone from the scorpion <i>Buthotus judacis</i> . Aug 20 1990 |
| Alzheimer's disease: a dissection of its etiology. May 1997 | | 14/6/111 10482606 PMID: 1365703 Molecular biology of the amyloid of Alzheimer's disease. An overview. 1992 | 14/6/132 09022839 PMID: 2357597 Domestic acid-induced neurovisceral toxic syndrome: characterization of an animal model and putative antibodies. May 1990 |
| 14/6/90 11895926 PMID: 9174065 | | 14/6/112 10372249 PMID: 7504355 Physiological production of the beta-amyloid protein and the mechanism of Alzheimer's disease. Oct 1993 | 14/6/133 08652506 PMID: 2742585 A N-methyltransferase in human brain catalyzes N-methylation of 1,2,3,4-tetrahydroisoquinoline into N-methyl-1,2,3,4-tetrahydroisoquinoline : a precursor of a dopaminergic neurotoxin, N-methylisiquinolium ion. Jun 30 1989 |
| Protection against quinolinic acid-mediated excitotoxicity in nigrostriatal dopaminergic neurons by endogenous kynurenic acid. Jun 1997 | | 14/6/113 10372248 PMID: 7504354 Genetic and molecular advances in Alzheimer's disease. Oct 1993 | 14/6/134 08403132 PMID: 2903469 Non-competitive antagonists of N-methyl-D-aspartate receptors protect cortical and hippocampal cell cultures against glutamate neurotoxicity. Aug 31 1988 |
| 14/6/91 11863477 PMID: 9135095 | | 14/6/114 10360381 PMID: 8239285 Transgenic animal models for Alzheimer's disease. Sep 24 1993 | 14/6/135 08235036 PMID: 3357353 Role of the aziridinium moiety in the in vivo cholinotoxicity of ethylcholine aziridinium ion (AβGA). Mar 1988 |
| Amyloid precursor proteins protect neurons of transgenic mice against acute and chronic excitotoxic injuries in vivo. May 1997 | | 14/6/115 10295842 PMID: 8104320 Steroid hormones protect spinal cord neurons from glutamate toxicity. Jul 1993 | 14/6/136 07934646 PMID: 3496809 A fluorometric determination of N-methyl-4-phenylpyridinium ion, using high-performance liquid chromatography. May 1 1987 |
| 14/6/92 11704108 PMID: 9086490 | | 14/6/116 10254388 PMID: 8393864 Xenoxins, a family of peptides from dorsal gland secretion of <i>Xenopus laevis</i> related to snake venom cytotoxins and neurotoxins. Aug 5 1993 | 16/6/4 15004869 PMID: 14529727 Intramuscular injection of 125I-botulinum neurotoxin-complex versus 125I-botulinum-free neurotoxin: time course of tissue distribution. Oct 2003 |
| Schwann cells as targets for neurotoxicants. Fall-Winter 1996 | | 14/6/117 10235995 PMID: 7687284 Peptide growth factors protect against ischemia in culture by preventing nitric oxide toxicity. Jul 1993 | |
| 14/6/93 11684569 PMID: 9013394 | | 14/6/118 10201828 PMID: 8510827 Amyloid beta protein precursor accumulates in swollen neuritis throughout rat brain with aging. Apr 16 1993 | |
| Stereospecific occurrence of a parkinsonism-inducing catechol isoquinoline, N-methyl(R)isaboline, in the human intraventricular fluid. 1996 | | 14/6/119 10074904 PMID: 8431601 Depressant insect selective neurotoxins from scorpion venom: chemistry, action, and gene cloning. 1993 | |
| 14/6/94 11530972 PMID: 8843103 | | | |
| A novel enzyme enantio-selectively synthesizes (R)isaboline, a precursor of a dopaminergic neurotoxin, N-methyl(R)isaboline. Jul 19 1996 | | | |
| 14/6/95 11458227 PMID: 8720122 | | | |
| Correlation among secondary structure, amyloid precursor protein accumulation, and neurotoxicity of amyloid beta(25-35) peptide as analyzed by single alanine substitution. Dec 1995 | | | |
| 14/6/96 11451990 PMID: 8752124 | | | |
| Neurotoxicity of a carboxy-terminal fragment of the Alzheimer's amyloid precursor protein. Sep 1996 | | | |
| 14/6/97 11438309 PMID: 8719038 | | | |
| Ion channel/hypothesis for Alzheimer amyloid peptide neurotoxicity. Oct 1995 | | | |
| 14/6/98 11397053 PMID: 8731376 | | | |
| Cytotoxicity of prion protein peptide (PrP106-126) differs in mechanism from the cytotoxic activity of the Alzheimer's disease amyloid peptide, A beta 25-35. Mar 1996 | | | |
| 14/6/99 11378444 PMID: 8663372 | | | |

- 16/6/2 14621936 PMID: 12381720
Evaluation of the therapeutic usefulness of botulinum neurotoxin B, C1, E, and F compared with the long lasting type A. Basis for distinct durations of inhibition of exocytosis in central neurons. Jan 10 2003
- 16/6/3 14269829 PMID: 12077369
Ca2+-induced changes in SNAREs and synaptotagmin I correlate with triggered exocytosis from chromaffin cells: insights gleaned into the signal transduction using trypsin and botulinum toxins. Jul 1 2002
- 16/6/4 13529067 PMID: 10497159
Protein kinase B stimulates the translocation of GLUT4 but not GLUT1 or transferin receptors in 3T3-L1 adipocytes by a pathway involving SNAP-23, synaptobrevin-2, and/or cellubrevin. Oct 1 1999
- 16/6/5 12657424 PMID: 10800941
A late phase of exocytosis from synaptosomes induced by elevated [Ca2+]i is not blocked by Cbstidial neurotoxins. May 2000
- 16/6/6 12360595 PMID: 9670008
Vesicle exocytosis stimulated by alpha-latrotoxin is mediated by atrophilin and requires both external and stored Ca2+. Jul 15 1998
- 16/6/7 11876880 PMID: 9153412
Botulinum neurotoxin B inhibits insulin-stimulated glucose uptake into 3T3-L1 adipocytes and cleaves cellubrevin unlike type A toxin which failed to proteolyze the SNAP-23 present. May 13 1997
- 16/6/8 11818598 PMID: 9115981
Importance of two adjacent C-terminal sequences of SNAP-25 in exocytosis from intact and permeabilized chromaffin cells revealed by inhibition with botulinum neurotoxins A and E. Mar 18 1997
- 16/6/9 11400081 PMID: 8723765
Productive and non-productive binding of botulinum neurotoxin A to motor nerve endings are distinguished by its heavy chain. May 1 1996
- 16/6/10 11371648 PMID: 8665909
Distinct exocytotic responses of intact and permeabilised chromaffin cells after cleavage of the 25-kDa synaptosomal-associated protein (SNAP-25) or synaptobrevin by botulinum toxin A or B. Mar 15 1996
- 16/6/11 11283823 PMID: 8611567
Botulinum neurotoxin C1 cleaves both syntaxin and SNAP-25 in intact and permeabilized chromaffin cells: correlation with its blockade of catecholamine release. Feb 27 1996
- 16/6/12 11179774 PMID: 7578132
Expression and purification of the light chain of botulinum neurotoxin A: a single mutation abolishes its cleavage of SNAP-25 and neurotoxicity after reconstitution with the heavy chain. Nov 21 1995
- 16/6/13 10946489 PMID: 7727408
Blockade by botulinum neurotoxin B of catecholamine release from adrenochromaffin cells correlates with its cleavage of synaptobrevin and a homologue present on the granules. Apr 25 1995
- 16/6/14 10806627 PMID: 7803399
Differences in the protease activities of tetanus and botulinum B toxins revealed by the cleavage of vesicle-associated membrane protein and various sized fragments. Dec 27 1994
- 16/6/15 10631531 PMID: 8020471
Botulinum A and the light chain of tetanus toxins inhibit distinct stages of Mg-ATP-dependent catecholamine exocytosis from permeabilised chromaffin cells. Jun 1 1994
- 16/6/16 10520769 PMID: 7508383
Antagonism of the intracellular action of botulinum neurotoxin type A with monoclonal antibodies that map to light-chain epitopes. Jan 15 1994
- 16/6/17 10368345 PMID: 8245989
- Botulinum A like type B and tetanus toxins fulfil criteria for being a zinc-dependent protease. Dec 1993
- 16/6/18 10315574 PMID: 8104936
A role for the interchain disulfide or its participating thiols in the internalization of botulinum neurotoxin A revealed by a toxin derivative that binds to ecto-acceptors and inhibits transmitter release intracellularly. Oct 5 1993
- 16/6/19 09931419 PMID: 1356988
Tetanus toxin and botulinum toxins type A and B inhibit glutamate, gamma-aminobutyric acid, aspartate, and met-enkephalin release from synaptosomes. Clues to the locus of action. Oct 25 1992
- 16/6/20 09811086 PMID: 1608558
Differences in the temperature dependencies of uptake of botulinum and tetanus toxins in Aplysia neurons. May 25 1992
- 16/6/21 09398634 PMID: 1674511
Heterologous combinations of heavy and light chains from botulinum neurotoxin A and tetanus toxin inhibit neurotransmitter release in Aplysia. May 25 1991
- 16/6/22 09340544 PMID: 1847184
Microtubule-disrupting drugs and A23187 reveal differences in the inhibition of synaptosomal transmitter release by botulinum neurotoxins types A and B. Mar 1991
- 16/6/23 09256990 PMID: 1981953
ADP-ribosylation of cerebrocortical synaptosomal proteins by cholera, pertussis and botulinum toxins. 1990
- 16/6/24 09255753 PMID: 1981913
Inhibition of neurotransmitter release by botulinum neurotoxins and tetanus toxin at Aplysia synapses: role of the constituent chains. 1990
- 16/6/25 09247901 PMID: 1963641
Clues to the multi-phasic inhibitory action of botulinum neurotoxins on release of transmitters. 1990
- 16/6/26 09209771 PMID: 2176612
Light chain of botulinum neurotoxin is active in mammalian motor nerve terminals when delivered via liposomes. Dec 17 1990
- 16/6/27 08995165 PMID: 2188647
Botulinum type F neurotoxin. Large-scale purification and characterization of its binding to rat cerebrocortical synaptosomes. May 15 1990
- 16/6/28 08918304 PMID: 2311761
Ca2(+)-dependent noradrenaline release from permeabilised PC12 cells is blocked by botulinum neurotoxin A or its light chain. Feb 26 1990
- 16/6/29 08831769 PMID: 2574722
Multiple domains of botulinum neurotoxin contribute to its inhibition of transmitter release in Aplysia neurons. Dec 25 1989
- 16/6/30 08771643 PMID: 2572418
Inhibition of transmitter release by botulinum neurotoxin A. Contribution of various fragments to the intoxication process. Oct 20 1989
- 16/6/31 08723105 PMID: 2550027
Effects of botulinum neurotoxin and Lambert-Eaton myasthenic syndrome IgG at mouse nerve terminals. 1989
- 16/6/32 08426206 PMID: 3197726
Involvement of the constituent chains of botulinum neurotoxins A and B in the blockade of neurotransmitter release. Nov 15 1988
- 16/6/33 08338446 PMID: 2900674
[Ganglionic synapses of Aplysia as a model for the study of the mechanism of action of botulinum neurotoxins] Les synapses ganglionnaires d'Aplysia comme modeles d'etude du mode d'action des neurotoxines botuliques. 1988
- 16/6/34 08271005 PMID: 2897693
Neurotransmitter release is blocked intracellularly by botulinum neurotoxin, and this requires uptake of both toxin polypeptides by a process mediated by the larger chain. Jun 1988
- 16/6/35 08267108 PMID: 2897427
Characterization of the inhibitory action of botulinum neurotoxin type A on the release of several transmitters from rat cerebrocortical synaptosomes. Jun 1988
- 16/6/36 08114019 PMID: 2830561
Selective location of acceptors for botulinum neurotoxin A in the central and peripheral nervous systems. Nov 1987
- 16/6/37 08053944 PMID: 3681717
The effects of in vitro application of purified botulinum neurotoxin at mouse motor nerve terminals. May 1987
- 16/6/38 07912843 PMID: 2439334
Botulinum toxin A blocks glutamate exocytosis from guinea-pig cerebral cortical synaptosomes. Jun 15 1987
- 16/6/39 07624489 PMID: 3015983
Interaction of 125I-labelled botulinum neurotoxins with nerve terminals. II. Autoradiographic evidence for its uptake into motor nerves by acceptor-mediated endocytosis. Aug 1986
- 16/6/40 07624488 PMID: 3733877
Interaction of 125I-labelled botulinum neurotoxins with nerve terminals. I. Ultrastructural autoradiographic localization and quantitation of distinct membrane acceptors for types A and B on motor nerves. Aug 1986
- 16/6/41 07555949 PMID: 2871643
Actions of beta-bungarotoxin on spontaneous release of transmitter at muscle end-plates treated with botulinum toxin. 1986
- 16/6/42 07455941 PMID: 3753681
Botulinum neurotoxin type B. Its purification, radioiodination and interaction with rat-brain synaptosomal membranes. Jan 15 1986
- 16/6/43 07316180 PMID: 3895579
A sensitive and useful radioimmunoassay for neurotoxin and its haemagglutinin complex from Clostridium botulinum. 1985
- 16/6/44 07204558 PMID: 6152294
Botulinum neurotoxin and dendrotoxin as probes for studies on transmitter release. 1984
- 16/6/45 06894193 PMID: 6694738
Acceptors for botulinum neurotoxin reside on motor nerve terminals and mediate its internalization. Feb 2-8 1984
- 16/6/46 06565837 PMID: 6339240
Radioiodination of botulinum neurotoxin type A with retention of biological activity and its binding to brain synaptosomes. Mar 15 1983
- 16/6/47 06370781 PMID: 6124898
The effects of purified botulinum neurotoxin type A on cholinergic, adrenergic and non-adrenergic, atropine-resistant autonomic neuromuscular transmission. Apr 1982
- 16/6/48 06325550 PMID: 7043781
Localization of sites for 125I-labelled botulinum neurotoxin at murine neuromuscular junction and its binding to rat brain synaptosomes. 1982
- 19/7/1 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.
11179774 PMID: 7578132
Expression and purification of the light chain of botulinum neurotoxin A: a single mutation abolishes its cleavage of SNAP-25 and neurotoxicity after reconstitution with the heavy chain.
Zhou L, de Paiva A, Liu D, Aoki R, Dolly J O
Department of Biochemistry, Imperial College of Science, Technology and Medicine, London.
Biochemistry (UNITED STATES) Nov 21 1995, 34 (46) p15175-81, ISSN 0006-2960

Journal Code: 0370623
 Publishing Model Print Document type: Journal Article Languages: ENGLISH Main
 Citation Owner: NLM
 Record type: MEDLINE; Completed
 Botulinum neurotoxin type A (BoNT/A) selectively and irreversibly inhibits acetylcholine release from peripheral nerve endings. While the toxin's heavy (H) chain contributes to neuronal binding and internalization, its light (L) chain is a Zn(2+)-dependent endoprotease that intracellularly cleaves synaptosomal-associated protein of Mr = 25 kDa (SNAP-25). For research and clinical exploitation of this uniquely-acting neurotoxin, recombinant wild-type L chain was produced together with a mutant in which His227 in the Zn(2+)-binding motif was substituted by Tyr. The PCR-amplified wild-type and mutant L chain genes were cloned, fused to the gene for maltose-binding protein, and expressed at high levels in *Escherichia coli*. The soluble fusion proteins were purified using amylose affinity chromatography, and, after factor Xa cleavage, the free L chains were isolated. The wild-type was shown to proteolyze SNAP-25 at a rate approaching that of the native chain while the mutant was inactive. Reconstitution of the pure wild-type L chain with native homogeneous H chain yielded a disulfide-linked dimer form that inhibited neuromuscular transmission *in vitro* and produced the symptoms of botulism *in vivo*. After reconstitution with the H chain, the Tyr227 mutant L chain failed to show any neuromuscular activity in either of these assays. This methodology allows, for the first time, routine preparation of recombinant forms of the L chain that are needed to decipher the molecular details of its interaction with substrate and, thereby, assist the design of effective inhibitors (ABSTRACT TRUNCATED AT 250 WORDS)
 Record Date Created: 19951226 Record Date Completed: 19951226

19/7/2 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.
 09931419 PMID: 1356988

Tetanus toxin and botulinum toxins type A and B inhibit glutamate, gamma-aminobutyric acid, aspartate, and mel-enkephalin release from synaptosomes. Clues to the locus of action.

McMahon H T; Foran P; Dolly J O; Verhage M; Wiegant V M; Nicholls D G
 Department of Biochemistry, University of Dundee, United Kingdom.
 Journal of biological chemistry (UNITED STATES) Oct 25 1992; 267 (30) p21338-43, ISSN 0021-9258 Journal Code: 2985121R Publishing Model Print Document type: Journal Article Languages: ENGLISH

Main Citation Owner: NLM Record type: MEDLINE; Completed
 Tetanus toxin (100 nM) when preincubated with guinea pig cerebrocortical synaptosomes for 45 min reduces the final extent of the KCl-evoked, Ca(2+)-dependent, glutamate transmitter release to 30% of non-intoxicated controls. Similarly, 100 nM Botulinum neurotoxins, types A and B, preincubated for 90 min inhibit release to 45-60% of non-intoxicated controls. The toxins preferentially attenuate a slow phase of KCl-evoked glutamate release which may be associated with synaptic vesicle mobilization. Tetanus toxin additionally inhibits the release of aspartate, gamma-aminobutyric acid and mel-enkephalin from the same preparation. Since amino acids and neuropeptides are released by distinct mechanisms, this indicates that the toxin affects a step common to both exocytotic pathways. When Ba2+ (which does not interact with calmodulin) is substituted for Ca2+, the control KCl-evoked release of each transmitter is unaffected and tetanus toxin is still inhibitory. Taken together these results implicate a calmodulin-independent locus (or loci) of action common to small- and large-dense-core vesicles and associated with vesicle transport.

Record Date Created: 19921125 Record Date Completed: 19921125

File 155:MEDLINE(R) 1951-2005/Sep 19 (c) format only 2005 Dialog

| Set | Items | Description |
|-----|--------|---|
| S1 | 6955 | DC=B3.300.390.400.200.' (CLOSTRIDIUM) |
| S2 | 111321 | DC=D12.776.828.' (RECOMBINANT PROTEINS) |
| S3 | 193 | S1 AND S2 |
| S4 | 7486 | DC=D24.185.926.640.' (NEUROTOXINS) |
| S5 | 4423 | DC=D24.185.926.123.179.' (BOTULINUM TOXINS) |
| S6 | 90 | S2 AND S5 |
| S7 | 37269 | 'MUTAGENESIS, SITE-DIRECTED' |
| S8 | 18657 | DC=G5.600.' (MUTAGENESIS) |
| S9 | 28 | S5 AND S7 |
| S10 | 5 | S5 AND S8 NOT S9 |
| S11 | 13 | S5 AND PRECURSOR |
| S12 | 1771 | DC=D24.185.926.123.893.' (TETANUS TOXIN) |
| S13 | 11 | S7 AND S12 |
| S14 | 13 | S12 AND PRECURSOR |
| S15 | 4 | S8 AND S12 NOT S13 |

13/6/11 14641734 PMID: 12553363
Tetanus toxin abolishes exocytosis of ROMK1 induced by inhibition of protein tyrosine kinase. Mar 2003

13/6/2 14483514 PMID: 12427026
Identification of a binding site for ganglioside on the receptor binding domain of tetanus toxin. Nov 19 2002

13/6/3 14143115 PMID: 11821389
Auto-ADP-ribosylation of *Pseudomonas aeruginosa* ExoS. Apr 5 2002

13/6/4 13946422 PMID: 11700044
Site-directed mutagenesis identifies active-site residues of the light chain of botulinum neurotoxin type A. Nov 16 2001

13/6/5 13829370 PMID: 11402045
Recombinant forms of tetanus toxin engineered for examining and exploiting neuronal trafficking pathways. Aug 17 2001

13/6/6 13645078 PMID: 11278003
Tyrosine-1290 of tetanus neurotoxin plays a key role in its binding to gangliosides and functional binding to neurones. Mar 23 2001

13/6/7 12853467 PMID: 10795737
Control of antigen presentation by a single protease cleavage site. Apr 2000

13/6/8 12245186 PMID: 9554265
Adjuvant effect of non-toxic mutants of *E. coli* heat-labile enterotoxin following intranasal, oral and intravaginal immunization. 1998

13/6/9 10722987 PMID: 7929330
Cytotoxic effects of a chimeric protein consisting of tetanus toxin light chain and anthrax toxin lethal factor in non-neuronal cells. Oct 21 1994

13/6/10 10607472 PMID: 7911329
A single mutation in the recombinant light chain of tetanus toxin abolishes its proteolytic activity and removes the toxicity seen after reconstitution with native heavy chain. Jun 7 1994

13/6/11 10186582 PMID: 8500613
Production of biologically active light chain of tetanus toxin in *Escherichia coli*. Evidence for the importance of the C-terminal 16 amino acids for full biological activity. Jun 1 1993

13/7/71 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.
12853467 PMID: 10795737

Control of antigen presentation by a single protease cleavage site.
Antoniou A N; Blackwood S L; Mazzeo D; Watts C
Department of Biochemistry, Wellcome Trust Biocentre, University of Dundee, United Kingdom.
Immunology (UNITED STATES) Apr 2000, 12 (4) p391-8, ISSN 1074-7613 Journal Code: 9432918
Publishing Model Print Document type: Journal Article Languages: ENGLISH
Main Citation Owner: NLM Record type: MEDLINE; Completed

Protein antigens require limited proteolytic processing to generate peptides for binding to class II MHC molecules, but the proteases and processing sites involved are largely unknown. Here we analyze the effect of eliminating the three major asparagine endopeptidase (AEP)-processing sites in the microbial antigen tetanus toxin C fragment. The mutant antigen is highly resistant to proteolysis by AEP and crude lysosomal extracts and is dramatically impaired in its ability to be processed and presented to T cells. Remarkably, processing at a single asparagine residue (1219) is obligatory for optimal presentation of many T cell epitopes in this antigen. These studies demonstrate that cleavage at a single processing site can be crucial for effective antigen presentation.

Record Date Created: 20000523 Record Date Completed: 20000523

13/7/9 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.
10722987 PMID: 7929330

Cytotoxic effects of a chimeric protein consisting of tetanus toxin light chain and anthrax toxin lethal factor in non-neuronal cells.

Arora N; Williamson L C; Leppla S H; Halpern J L
Laboratory of Microbial Ecology, NIDR, National Institutes of Health, Bethesda, Maryland 20892

Journal of biological chemistry (UNITED STATES) Oct 21 1994, 269 (42) p26165-71, ISSN 0021-9258 Journal Code: 2985121R Publishing Model Print Document type: Journal Article Languages: ENGLISH
Main Citation Owner: NLM Record type: MEDLINE; Completed

The light chain of tetanus toxin is a zinc endoprotease that inhibits neurotransmitter release by selective proteolysis of the synaptic vesicle-associated protein synaptobrevin/Vesicle-associated membrane protein. Cellubrevin is a homologue of synaptobrevin that is found in most cell types and is also a substrate for tetanus toxin. The lack of receptors for tetanus toxin on most cell types has made studies of tetanus toxin action in non-neuronal cells difficult. To characterize tetanus toxin effects in non-neuronal cells, a fusion protein consisting of the 254 amino-terminal amino acids of lethal factor (LF) of anthrax toxin and tetanus toxin light chain (LC) was prepared. This protein (LF-LC) inhibited evoked glycine release from primary spinal cord neurons at concentrations between 1.0 and 100 ng/ml. LF-LC was cytotoxic to RAW 264.7, ANA-1 cells (mouse macrophage cell lines), and Chinese hamster ovary cells in a dose-dependent manner. These effects required the presence of protective antigen, the receptor binding component of anthrax toxin. In contrast, LF-LC was not cytotoxic to RBL-2H3, Vero, or mouse hybridoma cell lines. Mutagenesis of conserved amino acids (His237 and Glu234) in the zinc-binding motif of LC resulted in fusion proteins having no biological activity. LF-LC did not inhibit regulated secretion of serotonin in RBL-2H3 cells or constitutive secretion in any non-neuronal cell lines as measured in several different assays. We suggest that the cytotoxic effects of LF-LC result from inhibition of a specific intracellular membrane fusion event mediated by cellubrevin.

Record Date Created: 19941122 Record Date Completed: 19941122

13/7/10 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.
10607472 PMID: 7911329

A single mutation in the recombinant light chain of tetanus toxin abolishes its proteolytic activity and removes the toxicity seen after reconstitution with native heavy chain.

Li Y; Foran P; Fairweather N F; de Paiva A; Weller U; Dougan G; Dolly J O
Department of Biochemistry, Imperial College, London, U.K.

Biochemistry (UNITED STATES) Jun 7 1994, 33 (22) p7014-20, ISSN 0006-2960 Journal Code: 0370623
Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM

Record type: MEDLINE; Completed

Specific proteolysis by the tetanus toxin light chain of a vesicle-associated membrane protein (VAMP) involved in exocytosis is thought to underlie its intracellular blockade of neurotransmitter release. To substantiate this mechanism, recombinant light chain was expressed as a maltose binding protein-light chain fusion product in *Escherichia coli*. After purification of affinity chromatography and cleavage with factor Xa, the resultant light chain was isolated and its identity confirmed by Western blotting and N-terminal sequencing. It exhibited activity similar to that of the native light chain in proteolyzing its target in isolated bovine small synaptic vesicles and in hydrolyzing a 62-residue synthetic polypeptide spanning the cleavage site of the substrate. The importance of Glu234 in the catalytic activity of the light chain, possibly analogous to Glu143 of thermolysin, was examined using site-directed mutagenesis. Changing Glu234 to Ala abolished the protease activity of the light chain, but its ability to bind the polypeptide substrate was retained. Each recombinant light chain could be reconstituted with the heavy chain of tetanus toxin, yielding the same level of disulfide-linked species as the two native chains. Whereas the toxin formed with wild-type light chain exhibited appreciable neuromuscular paralysis activity and mouse lethality, the equivalent dichain material containing the Ala234 mutant lacked neurotoxicity in both the *in vitro* and *in vivo* assays. Thus, these results demonstrate directly, for the first time, that the lethality of tetanus toxin and its inhibition of exocytosis in intact neurons are attributable largely, if not exclusively, to endoprotease activity.

Record Date Created: 19940708 Record Date Completed: 19940708

13/7/11 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.
10186582 PMID: 8500613

Production of biologically active light chain of tetanus toxin in *Escherichia coli*. Evidence for the importance of the C-terminal 16 amino acids for full biological activity.

Fairweather N.F., Sanders D., Slater D., Hudel M., Habermann E., Weller U.
Department of Cell Biology, Wellcome Foundation Ltd., Beckenham, Kent, UK.
FEBS letters (NETHERLANDS) Jun 1 1993, 323 (3) p218-22, ISSN 0014-5793 Journal Code: 0155157
Publishing Model Print Document type: Journal Article Languages: ENGLISH Main Citation Owner: NLM
Record type: MEDLINE; Completed

The activity of the light (L) chain of tetanus toxin, and of mutants constructed by site-directed mutagenesis, was studied by expression and purification of the proteins from *E. coli*. Wild-type recombinant L chain (pTe187) was active in the inhibition of exocytosis from cultured bovine adrenal chromaffin cells, although at a level 5-15% of that of L chain purified from tetanus toxin. L chain mutants which terminated at Leu-438 (pTe189), or which contained a Cys-to-Ser mutation at residue 439 (pTe188) were equally as active as the full-length recombinant protein. The reduced activity of pTe187 L chain correlated with C-terminal proteolysis of the protein upon purification. A tryptic fragment derived from native light chain and which terminated at Leu-434 also showed reduced activity in the exocytosis assay, consistent with a requirement of the C-terminal region of the L chain for maximal activity. pTe187 L chain, but neither of the mutants, could be associated with purified H (heavy) chain to form a covalent dimer which induced the symptoms of tetanus in mice. The ability to form biologically active toxin using recombinant L chain will be of great value in structure-function studies of tetanus toxin.
Record Date Created: 19930629 Record Date Completed: 19930629

14/6/1 14048520 PMID: 11814298
Characterization of tetanus toxin, neat and in culture supernatant, by electrospray mass spectrometry. Feb 15 2002

14/6/2 13765559 PMID: 11427819
[Mechanism of action and therapeutic uses of botulinum and tetanus neurotoxins] Mécanismes d'action et utilisations thérapeutiques des neurotoxines botuliques et tétaniques. May 2001

14/6/3 10753456 PMID: 7957894
Fusion complex formation protects synaptobrevin against proteolysis by tetanus toxin light chain. Oct 24 1994

14/6/4 10357199 PMID: 7901925
Tetanus and botulinum neurotoxins: a new group of zinc proteases. Sep 1993

14/6/5 09247900 PMID: 2074546
Chains and fragments of tetanus toxin, and their contribution to toxicity. 1990

14/6/6 08817045 PMID: 24803377
Processing of tetanus toxin by human antigen-presenting cells. Evidence for donor and epitope-specific processing pathways. Dec 15 1989

14/6/7 08402736 PMID: 3054567 Record Identifier: 89040247
Tetanus toxin: biochemical and pharmacological comparison between its protoxin and some isotoxins obtained by limited proteolysis. Aug 1988

14/6/8 07427497 PMID: 3000624
Neural induction and in vitro initial expression of neurofilament and tetanus toxin binding site moieties in amphibians. Jan 1986

14/6/9 07333881 PMID: 2863321
In vitro differentiation of neuronal precursor cells from amphibian late gastrulae: morphological, immunocytochemical studies, biosynthesis, accumulation and uptake of neurotransmitters. Apr 1985

14/6/10 07304663 PMID: 2410447
Cell type specificity and developmental expression of the L2/HNK-1 epitopes in mouse cerebellum. Aug 1985

14/6/11 06798577 PMID: 6653877
Neuronal acquisition of tetanus toxin binding sites: relationship with the fast mitotic cycle. Dec 1983

14/6/12 06720680 PMID: 6194025
Tracing the development of oligodendrocytes from precursor cells using monoclonal antibodies, fluorescence-activated cell sorting, and cell culture. Nov 1983

14/6/13 06043343 PMID: 7015184
Tetanus toxin: a marker of amphibian neuronal differentiation in vitro. Mar 10 1981

14/5/7 DIALOG(R)File 155:MEDLINE(R) (c) format only 2005 Dialog. All rts. reserv.
08402736 PMID: 3054567 Record Identifier: 89040247
Tetanus toxin: biochemical and pharmacological comparison between its protoxin and some isotoxins obtained by limited proteolysis.
Weller U., Mauler F., Habermann E.
Rudolf-Buchheim-Institut für Pharmakologie, Justus-Liebig-Universität Gießen, Federal Republic of Germany.
Naunyn-Schmiedeberg's archives of pharmacology (GERMANY, WEST) Aug 1988, 338 (2) p99-106, ISSN 0028-1298
Journal Code: 0326264 Publishing Model Print Document type: Journal Article Languages: ENGLISH

Main Citation Owner: NLM Other Citation Owner: NASA Record type: MEDLINE; Completed Subfile: INDEX MEDICUS
Single-chain tetanus toxin (toxin S) was prepared from short-term cultures by lysis under protection with protease inhibitors, precipitation with 40% ammonium sulfate, gel filtration, and chromatography on DEAE ion exchanger. Its limited proteolysis by trypsin, post-arginine cleaving enzyme from mouse submaxillary gland and doxipain led to bichainal derivatives (BT, BA, BCI) consisting of a heavy chain and a larger version of the light chain. The latter was then converted by trypsin into a small version which comigrated with the light chain of bichainal extracellular toxin (BE). The light chain produced by chymotrypsin (BC) and elastase (BE1) was of intermediate size. The nick region serves as substrate for all esteraseproteases investigated and comprises between one and two kDa. Limited proteolysis increased the hydrophilicity (BT greater than BE greater than S) in hydrophobic interaction HPLC, and anionic behaviour (BC greater than BE greater than BT greater than S) in DEAE ion exchanger HPLC. The bichainal toxins assessed (BC, BE or BT) were about two times more toxic than toxin S (LD50, mouse s.c. 2 ng/kg vs. 4 ng/kg). They were five to twelve times more potent than toxin S in three in vitro assays measuring the prevention of neurotransmitter release, i.e. on the phrenic nerve-hemidiaphragm preparation of the mouse (acetylcholine, with toxin BE and BT), on primary brain cell cultures from the mouse (3H-fluoradrenaline, with toxin BA, BC, BE and BT), and on brain homogenate from rats (3H-fluoradrenaline, with toxin BA, BC, BE and BT). Thus single-chain toxin is a less potent precursor of, or protoxin for, various bichainal isotoxins (ABSTRACT TRUNCATED AT 250 WORDS)
Tags: Comparative Study, In Vitro; Research Support, Non-U.S. Gov't
Descriptors: *Tetanus Toxin--isolation and purification--IP; Animals; Brain--drug effects--DE; Brain--metabolism--ME; Chromatography--methods--MT; Hydrolysis; Mice; Neuromuscular Junction--drug effects--DE; Norepinephrine--metabolism--ME; Peptide Fragments--isolation and purification--IP; Peptide Fragments--metabolism--ME; Peptide Fragments--toxicity--TO; Peptide Hydrolases; Rats; Tetanus Toxin--metabolism--ME; Tetanus Toxin--toxicity--TO
CAS Registry No.: 0 (Peptide Fragments); 0 (Tetanus Toxin); 51-41-2 (Norepinephrine)
Enzyme No.: EC 3.4.- (Peptide Hydrolases)
Record Date Created: 19881221 Record Date Completed: 19881221

15/6/1 14918480 PMID: 12764154
Activation of store-operated calcium channels: assessment of the role of snare-mediated vesicular transport. Aug 15 2003

15/6/2 14211526 PMID: 12010992
The LTR72 mutant of heat-labile enterotoxin of *Escherichia coli* enhances the ability of peptide antigens to elicit CD4(+) T cells and secrete gamma interferon after coapplication onto bare skin. Jun 2002

15/6/3 13668786 PMID: 11306125
Active-site mutagenesis of tetanus neurotoxin implicates TYR-375 and GLU-271 in metalloproteolytic activity. Aug 2001

15/6/4 13014512 PMID: 10972823
Analysis of mutants of tetanus toxin Hc fragment: ganglioside binding, cell binding and retrograde axonal transport properties. Sep 2000